

Routine Recommendations for Meningococcal Vaccines¹

Vaccine Type	Brand Names	Routine Recommendation for People Who Are Not in a Risk Group ^{1,2,3}	General Guidelines
MenACWY ²	Menactra® MenQuadfi® Menveo®	One dose at age 11-12 years; booster dose at age 16 years	<ul style="list-style-type: none"> • 1 dose required⁴ at 11 years of age or older upon entry into 7th grade or higher • Products are interchangeable if age appropriate, but same vaccine is recommended for complete series • Intramuscular injection
MenB ³	Bexsero® Trumenba®	Shared clinical decision making for persons 16 years through 23 years without high-risk conditions: <ul style="list-style-type: none"> • Bexsero: 2 dose series at least 1 month apart OR • Trumenba: 2 dose series at least 6 months apart 	<ul style="list-style-type: none"> • Products are not interchangeable • Intramuscular injection • MenB vaccine is not routinely recommended at 11-12 years of age

¹ For Child and Adolescent Immunization Schedule by Age at: www.cdc.gov/vaccines/schedules/hcp/imz/child-adolescent.html

² For more information regarding Meningococcal Conjugate Vaccine (MenACWY), including guidance for catch-up and for persons who are at high-risk, refer to the Meningococcal Conjugate Quick Look at: www.michigan.gov/vaccinequicklooks and Immunize.org's handout titled, "Meningococcal ACWY vaccine recommendations by Age and Risk Factor" at: www.immunize.org/catg.d/p2018.pdf

³ For more information regarding Meningococcal Serogroup B (MenB), including guidance for catch-up and for persons who are at high-risk, refer to the Meningococcal Serogroup B Quick Look at: www.michigan.gov/vaccinequicklooks and Immunize.org's handout titled, "Meningococcal B Vaccine Recommendations by Age and Risk Factor" at: www.immunize.org/catg.d/p2035.pdf

⁴ For vaccines required for school entry in Michigan, refer to "Vaccines Required for School Entry in Michigan" at: https://mcir.org/wp-content/uploads/2021/08/SchoolEntryReqVaccinesParentsMI_5.3.2021approvedfinalpublish.pdf